

# Quarks and baryons in QCD at finite density

Yu.A. Simonov, M.A.Trusov  
*ITEP, Moscow, Russia*

## Abstract

The mechanism of string creation for light quarks developed earlier is considered for nonzero quark chemical potential. A strong modification of the confining string due to finite quark density (chemical quark potential  $\mu$ ) is observed. As a surprising result in a multiquark system with a common string junction an attractive well appears of radius  $\mu/\sigma$  and of an average depth equal to  $\mu$ , which implies formation of multiquark hadrons. Preliminary estimates predict a new phase transition to multiquark hadron phase at rather high densities (in heavy ion collisions or in neutron stars) when neutron matter is compressed to 3-4 normal nuclear densities.

## 1 Introduction

Recently the high-density effects in QCD attracted a lot of attention because of the possible density phase transition [1, 2]. On the fundamental level the physical expectation of any phase transition may be connected to the possible reconstruction of the vacuum and for that one needs that the energy density to be of the order of the vacuum energy density  $\varepsilon_{cr} \sim \varepsilon_{vac} \approx -\frac{11}{3}N_c \frac{\alpha_s}{32\pi} \langle (F_{\mu\nu}^a)^2 \rangle$ ,  $\varepsilon_{cr} \sim 1 \text{ GeV/fm}^3$  which provides the drastic change in the vacuum structure and may cause the phase transition. In case of temperature phase transition the corresponding energy density  $\varepsilon$  is indeed of the order of  $\varepsilon_{cr}$ , and as it was first argued in [3] and later measured on the lattice [4], the QCD vacuum is strongly transformed in a way, that most part of colorelectric fields evaporate above  $T_c$ . In this way  $T_c$  was calculated through  $\varepsilon_{cr}$  and found in good agreement with lattice data [5], see [6] for a review. It was shown earlier in [7], that Chiral Symmetry Breaking (CSB) occurs due to confinement (more explicitly its colorelectric scalar components), which means that CSB should disappear simultaneously with confinement. In another way in the framework of hadron resonance gas and making use of the low-energy theorems at  $T \neq 0$  a similar conclusion was obtained through application of the effective dilaton Lagrangian to gluodynamics and to QCD [8].

The same type of arguments for the density phase transition would imply that at the baryon density of  $\sim 1 \text{ nucleon/fm}^3$ , i.e.  $3 \div 6$  times higher than the standard nuclear density, the vacuum can be reconstructed in such a way, that part of fields, e.g. the colorelectric fields responsible for confinement, disappear above critical density. The actual calculations in [5] have supported this conclusion and give critical chemical potential  $\mu_q(T=0) = 0.6 \text{ GeV}$ .

This is the picture of deconfinement due to high density. However, of much more practical importance can be possible phase transitions in the confining region, at  $2 \div 3$  normal nuclear densities.

It is the purpose of the present paper to start the investigation of the role of density on the vacuum fields in general and confinement in particular.

In this paper we ask ourselves a short and simple question: how the nonzero baryonic chemical potential  $\mu$  acts on the confinement of light quarks, and come to the unexpected answer, that the confining string of light quarks is destroyed gradually by  $\mu$  in such a way, that the one part of string, near the string junction is eaten by the nonzero  $\mu$ , while at distant  $r$  the string survives. We show that this can make multiquark states (MQS) more advantageous and thus cause a new type of phase transition. The plan of the paper is as follows. In section 2 the effective Lagrangian is derived from the QCD Lagrangian with nonzero  $\mu$  and their solution is discussed in section 3. Section 4 generalizes results to the case of baryons and multiquark bags. Section 5 is devoted to the physical implications of results and prospectives.

## 2 Derivation of Effective $4q$ Lagrangian

One starts with the QCD partition function in presence of quark chemical potential  $\mu$  in the Euclidean space-time, and we begin with the zero temperature,  $T = 0$ .

$$Z = \int DAD\psi D\psi^+ e^{-S_0(A) + \int {}^f\psi^+ (i\hat{\partial} + im - i\mu\gamma_4 + g\hat{A}) {}^f\psi d^4x} \quad (1)$$

where  $S_0(A) = \frac{1}{4} \int (F_{\mu\nu}^a(x))^2 d^4x$ ,  $m$  is the current quark mass (mass matrix  $\hat{m}$  in SU(3)), and the quark operator  ${}^f\psi_{a\alpha}(x)$  has flavor index  $a(f = 1, \dots, n_f)$ , color index  $a(a = 1, \dots, N_c)$  and Lorenz bispinor index  $\alpha(\alpha = 1, 2, 3, 4)$ , and we use the contour gauge [9] to express  $A_\mu(x)$  in terms of  $F_{\mu\nu}$ . One has for the contour  $z_\mu(s, x)$  starting at point  $x$  and ending at  $Y = z(0, x)$

$$A_\mu(x) = \int_0^1 ds \frac{\partial z_\nu(s, x)}{\partial s} \frac{\partial z_\rho(s, x)}{\partial x_\mu} F_{\nu\rho}(z(s)) \equiv \int_Y^x d\Gamma_{\mu\nu\rho}(z) F_{\nu\rho}(z). \quad (2)$$

Integrating out the gluonic fields  $A_\mu(x)$ , one obtains

$$Z = \int D\psi D\psi^+ e^{\int {}^f\psi^+ (i\hat{\partial} + im - i\mu\gamma_4) {}^f\psi d^4x} e^{L_{EQL}^{(2)} + L_{EQL}^{(3)} + \dots} \quad (3)$$

where the EQL proportional to  $\langle\langle A^n \rangle\rangle$  is denoted by  $L_{EQL}^{(n)}$ ,

$$L_{EQL}^{(2)} = \frac{g^2}{2} \int d^4x d^4y {}^f\psi_{a\alpha}^+(x) {}^f\psi_{b\beta}(x) {}^g\psi_{c\gamma}^+(y) {}^g\psi_{d\varepsilon}(y) \langle A_{ab}^{(\mu)}(x) A_{cd}^{(\nu)}(y) \rangle \gamma_{\alpha\beta}^{(\mu)} \gamma_{\gamma\varepsilon}^{(\nu)} \quad (4)$$

Average of gluonic fields can be computed using (2) as (see [10] for details of derivation)

$$g^2 \langle A_{ab}^{(\mu)}(x) A_{cd}^{(\nu)}(y) \rangle = \frac{\delta_{bc} \delta_{ad}}{N_c} \int_0^x du_i \alpha_\mu(u) \int_0^y dv_k \alpha_\nu(v) D^{(E,H)}(u-v) (\delta_{\mu\nu} \delta_{ik} - \delta_{i\nu} \delta_{k\mu}), \quad (5)$$

where  $D^{(E,H)}(x)$  is the correlator  $\langle E_i(x)E_i(0) \rangle$  or  $\langle H_i(x)H_i(0) \rangle$ . As it was argued in [10] the dominant contribution at large distances from the static antiquark is given by the color-electric fields, therefore at the first stage we shall write down explicitly  $L_{EQL}^{(2)}(el)$  for this case, i.e. taking  $\mu = \nu = 4$ . As a result one has[10]

$$L_{EQL}^{(2)}(el) = \frac{1}{2N_c} \int d^4x \int d^4y {}^f\psi_{a\alpha}^+(x) {}^f\psi_{b\beta}(x) {}^g\psi_{b\gamma}^+(y) {}^g\psi_{a\varepsilon}(y) \gamma_{\alpha\beta}^{(4)} \gamma_{\gamma\varepsilon}^{(4)} J^E(x, y) \quad (6)$$

where  $J^E(x, y)$  is

$$J^E(x, y) = \int_0^x du_i \int_0^y dv_i D^E(u - v), \quad i = 1, 2, 3. \quad (7)$$

One can form bilinears  $\Psi_{\alpha\varepsilon}^{fg} \equiv {}^f\psi_{a\alpha}^+ {}^g\psi_{a\varepsilon}$  and project using Fierz procedure given isospin and Lorentz structures,  $\Psi_{\alpha\varepsilon}^{fg} \rightarrow \Psi^{(n,k)}(x, y)$ . Here we consider only  $\psi^+\psi$  bosonization. With the help of the standard bosonization trick (here  $\tilde{J} \equiv \frac{1}{N_c} J^E$ )

$$e^{-\Psi\tilde{J}\Psi} = \int (\det \tilde{J})^{1/2} D\chi \exp[-\chi\tilde{J}\chi + i\Psi\tilde{J}\chi + i\chi\tilde{J}\Psi] \quad (8)$$

$$Z = \int D\psi D\psi^+ D\chi \exp L_{QML} \quad (9)$$

one obtains the effective Quark-Meson Lagrangian (QML)

$$L_{QML}^{(2)} = \int d^4x \int d^4y \left\{ {}^f\psi_{a\alpha}^+(x) [(i\hat{\partial} + im - i\mu\gamma_4)_{\alpha\beta} \delta(x - y) + iM_{\alpha\beta}^{(fg)}(x, y)] {}^g\psi_{a\beta}(y) - \frac{1}{N_c} \chi^{(n,k)}(x, y) J^E(x, y) \chi^{(n,k)}(y, x) \right\} \quad (10)$$

and the effective quark-mass operator is

$$M_{\alpha\beta}^{(fg)}(x, y) = \sum_{n,k} \chi^{(n,k)}(x, y) O_{\alpha\beta}^{(k)} t_{fg}^{(n)} \tilde{J}(x, y). \quad (11)$$

Here the operator  $\hat{O}$  is a set of all irreducible combinations of Dirac matrixes.

The QML in Eq.(10)  $L_{QML}^{(2)}$  contains functions  $\chi^{(n,k)}$  which are integrated out in (9), and the standard way is to find  $\chi^{(n,k)}$  from the stationary point of  $L_{QML}^{(2)}$ . Limiting oneself to the scalar and pseudoscalar fields and using the nonlinear parametrization one can write for the operator  $\hat{M}$  in (10)

$$\hat{M}(x, y) = M_S(x, y) \hat{U}(x, y), \quad \hat{U} = \exp(i\gamma_5 \hat{\phi}), \quad \hat{\phi}(x, y) = \phi^f(x, y) t^f. \quad (12)$$

After integrating out the quark fields one obtains the ECL in the form

$$L_{ECL}^{(2)}(M_S, \hat{\phi}) = -2n_f N_c (J^E(x, y))^{-1} M_S^2(x, y) + N_c \text{tr} \log[(i\hat{\partial} + im - i\mu\gamma_4) \hat{1} + iM_S \hat{U}]. \quad (13)$$

The stationary point equations  $\frac{\delta L_{ECL}^{(2)}}{\delta M_s} = \frac{\delta L_{ECL}^{(2)}}{\delta \hat{\phi}} = 0$  at  $\hat{\phi} = \hat{\phi}_0$ ,  $M_s = M_s^{(0)}$  immediately show that  $\hat{\phi}_0 = 0$  and  $M_s^{(0)}$  satisfies nonlinear equation

$$iM_S^{(0)}(x, y) = 4\text{tr} S J^E(x, y) = (\gamma_4 S \gamma_4) J^E(x, y), \quad S(x, y) = -[i\hat{\partial} + im - i\mu\gamma_4 + iM_S \hat{U}]_{x,y}^{-1}. \quad (14)$$

This equation plays the role of the gap equation and is the main point of our further investigation. For  $\mu = 0$  this was done in [10] and in the next section we find how results of [10] are modified by the nonzero  $\mu$ .

### 3 The confining string at nonzero $\mu$

Our basic equations (6),(7) are nonlocal in time because of the integral over  $dx_4 dy_4$  in (10). This nonlocality and the parameter which it governs can be handled most easily, when one uses instead of  $M(z, z')$ ,  $S(z, z')$  the Fourier transforms.

$$S(z_4 - z'_4, \mathbf{z}, \mathbf{z}') = \int e^{ip_4(z_4 - z'_4)} S(p_4, \mathbf{z}, \mathbf{z}') \frac{dp_4}{2\pi} \quad (15)$$

and the same for  $M(z, z')$ . Then from (14) one obtains a system of equations

$$(\hat{p}_4 - i\hat{\partial}_z - im + i\mu\gamma_4)S(p_4, \mathbf{z}, \mathbf{w}) - i \int M(p_4, \mathbf{z}, \mathbf{z}') S(p_4, \mathbf{z}', \mathbf{w}) d\mathbf{z}' = \delta^{(3)}(\mathbf{z} - \mathbf{w}) \quad (16)$$

To simplify matter, one assumes for  $D^E(x)$  the Gaussian form,  $D^E(x) = D(0) \exp\left(-\frac{x^2}{4T_g^2}\right)$ . Then for  $M(p_4, \mathbf{z}, \mathbf{w})$  one has

$$iM(p_4, \mathbf{z}, \mathbf{w}) = 2\sqrt{\pi}T_g \int \frac{dp'_4}{2\pi} e^{-(p_4 - p'_4)^2 T_g^2} \times \\ \times [J^E(\mathbf{z}, \mathbf{w}) \gamma_4 S(p'_4, \mathbf{z}, \mathbf{w}) \gamma_4] \quad (17)$$

where  $J^E$  is defined in (7) and we have factored out the time-dependent exponent, using the Gaussian representation of  $D(u)$ .

All dependence of  $M$  on  $p_4$  as can be seen in (17) is due to the factor  $\exp[-(p_4 - p'_4)^2 T_g^2]$  and disappears in the limit when  $T_g$  goes to zero, while the string tension  $\sigma \sim D(0)T_g^2$  is kept fixed. This limit can be called the string limit of QCD, and we shall study its consequences for equations (16),(17) in this section.

So in the string limit, with  $M$  independent of  $p_4$ , let us consider the Hermitian Hamiltonian

$$\hat{H}\psi_n \equiv \left(\frac{\alpha_i}{i} \frac{\partial}{\partial z_i} + \beta m - \mu\right) \psi_n(\mathbf{z}) + \beta \int M(p_4 = 0, \mathbf{z}, \mathbf{z}') \psi_n(\mathbf{z}') d^3 \mathbf{z}' = \tilde{\varepsilon}_n(\mu) \psi_n(\mathbf{z}) \quad (18)$$

with eigenfunctions  $\psi_n$  satisfying usual orthonormality condition

$$\int \psi_n^+(x) \psi_m(x) d^3 x = \delta_{nm},$$

From (18) it is clear, that one can redefine  $\tilde{\varepsilon}_n(\mu) + \mu \equiv \varepsilon_n$ , and  $\varepsilon_n$  and  $\psi_n$  do not depend on  $\mu$ . Therefore in all subsequent formulas one can use the same equations as in [10], but with the replacement  $\varepsilon_n \rightarrow \varepsilon_n - \mu$ . In particular, the Green's function  $S$  can be expressed as

$$S(p_4, \mathbf{x}, \mathbf{y}) = \sum_n \frac{\gamma_4 \psi_n(\mathbf{x}) \psi_n^+(\mathbf{y})}{p_4 - i(\varepsilon_n - \mu)} \quad (19)$$

Inserting (19) into (17) one has integrals of the type:

$$\int_{-\infty}^{\infty} \frac{dp'_4}{2\pi} \frac{\gamma_4 e^{-(p_4 - p'_4)^2 T_g^2}}{(p'_4 - i(\varepsilon_n - \mu))} = \frac{i}{2} \gamma_4 \text{sign}(\varepsilon_n - \mu) (1 + 0(p_4 T_g, |\varepsilon_n| T_g)) \quad (20)$$

We are thus led to the following expression for  $M$  in the string limit

$$M(p_4 = 0, \mathbf{z}, \mathbf{w}) = \sqrt{\pi} T_g J^E(\mathbf{z}, \mathbf{w}) \gamma_4 \Lambda(\mathbf{z}, \mathbf{w}) \quad (21)$$

where the definition is used

$$\Lambda(\mathbf{z}, \mathbf{w}) = \sum_n \psi_n(\mathbf{z}) \text{sign}(\varepsilon_n - \mu) \psi_n^+(\mathbf{w}) \quad (22)$$

Let us disregard for the moment the possible appearance in  $M$  of the vector component (proportional to  $\gamma_\mu, \mu = 1, 2, 3, 4$ ) and concentrate on the scalar contribution only, since that is responsible for CSB and confinement. Then one can look for solutions of the Dirac equation (18) in the following form [10]

$$\psi_n(\vec{r}) = \frac{1}{r} \begin{pmatrix} G_n(r) \Omega_{jlM} \\ i F_n(r) \Omega_{j'l'M} \end{pmatrix} \quad (23)$$

where  $l' = 2j - l$ , and introducing the parameter  $\kappa(j, l) = (j + \frac{1}{2}) \text{sign}(l - j)$ , and replacing  $M$  by a local operator (the generalization to the nonlocal case is straightforward but cumbersome, for a possible change in the nonlocal case see [10]), we obtain a system of equations

$$\begin{cases} \frac{dG_n}{dr} + \frac{\kappa}{r} G_n - (\varepsilon_n - \mu + m + M_{scal}(r) - M_{vect.}(r)) F_n = 0 \\ \frac{dF_n}{dr} - \frac{\kappa}{r} F_n + (\varepsilon_n - \mu - m - M_{scal}(r) - M_{vect.}(r)) G_n = 0 \end{cases} \quad (24)$$

where we suggest  $M = M_{scal} + \gamma_4 M_{vect.}$

Eq.(23) possesses a symmetry  $(\varepsilon_n - \mu, G_n, F_n, \kappa) \leftrightarrow (\mu - \varepsilon_n, F_n, G_n, -\kappa)$  which means that for any solution of the form (22) corresponding to the eigenvalue  $\varepsilon_n - \mu$ , there is another solution of the form

$$\psi_{\mu - \varepsilon_n}(r) = \frac{1}{r} \begin{pmatrix} F_n(r) \Omega_{j'l'M} \\ i G_n(r) \Omega_{jlM} \end{pmatrix} \quad (25)$$

corresponding to the eigenvalue  $(\mu - \varepsilon_n)$ .

Therefore the difference, which enters (22) can be computed in terms of  $F_n, G_n$  as follows

$$\Lambda(\mathbf{z}, \mathbf{w}) = \Lambda_0(\mathbf{z}, \mathbf{w}) - \Delta\Lambda(\mathbf{z}, \mathbf{w}) \quad (26)$$

where  $\Lambda_0$  is the value of  $\Lambda$  for  $\mu = 0$ , i.e. the same as in [10], while  $\Delta\Lambda$  is defined as

$$\Delta\Lambda(\mathbf{z}, \mathbf{w}) = 2 \sum_{0 < \varepsilon_n < \mu} \psi_n(\mathbf{z}) \psi_n^+(\mathbf{w}). \quad (27)$$

Using decomposition (23) one can write  $\Delta\Lambda$  as

$$\Delta\Lambda(\mathbf{r}, \mathbf{r}') = 2 \sum_{0 < \varepsilon_n, \mu} \begin{pmatrix} G_n G_n^+ \Omega \Omega^+, & -i G_n F_n^+ \Omega \Omega'^+ \\ i F_n G_n^+ \Omega' \Omega^+, & F_n F_n^+ \Omega' \Omega'^+ \end{pmatrix} \quad (28)$$

where we have denoted  $\Omega \equiv \Omega_{jlM}, \Omega' = \Omega_{j'l'M}$ , and we disregard nondiagonal part of  $\Lambda$ .

At this point one can follow the relativistic WKB method for Dirac equation [11] applied to calculation of  $\Lambda$  in [10] in case of  $\mu = 0$ . The classically available region for

$\psi_n(\mathbf{r})$  with energy  $\varepsilon_n \equiv \varepsilon$  is  $r_{\min} \leq r \leq r_{\max}$ , where  $r_{\max, \min} = \frac{\varepsilon^2 \pm \sqrt{\varepsilon^2 - 4\sigma^2 \kappa^2}}{2\sigma^2}$ , and the summation over  $n$  in (27) transforms into integration over  $d\varepsilon$ , with the lower limit (for a given  $r$ )  $\varepsilon_{\min} = \sigma r$ . In this way one has for the upper diagonal element in (28).

$$\Delta\Lambda(+, +) = \frac{2\sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \int_1^{\mu/\sigma r} d\tau \frac{\tau + 1}{\sqrt{\tau^2 - 1}} \cos(a\sqrt{\tau^2 - 1}) \theta(\mu - \sigma r) \quad (29)$$

with  $a = \sigma r|r - r'|$ , and we keep  $r \approx r'$  everywhere except for  $a$ , since for large  $a$  (when  $r$  is far from  $r'$ ) both  $\Lambda$  and  $\Delta\Lambda$  fast decrease.

In a similar way for the lower diagonal element in (28) one has

$$\Delta\Lambda(-, -) = \frac{2\sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \int_1^{\mu/\sigma r} d\tau \frac{\tau - 1}{\sqrt{\tau^2 - 1}} \cos(a\sqrt{\tau^2 - 1}) \theta(\mu - \sigma r) \quad (30)$$

and taking  $\Lambda_0$  in (25) from [10] the resulting form for  $\Lambda$  (26) is

$$\begin{aligned} \Lambda(\mathbf{r}, \mathbf{r}') \equiv \beta \Lambda_{\text{scalar}} + \hat{1} \Lambda_{\text{vector}} &= \frac{\beta \sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \int_{\mu/\sigma r}^{\infty} \frac{d\tau \cos(a\sqrt{\tau^2 - 1})}{\sqrt{\tau^2 - 1}} - \\ &- \hat{1} \frac{2\sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \int_1^{\mu/\sigma r} \frac{\tau d\tau}{\sqrt{\tau^2 - 1}} \cos(a\sqrt{\tau^2 - 1}). \end{aligned} \quad (31)$$

Here  $\beta \equiv \gamma_4$ , and  $\hat{1}$  is the unit Dirac matrix, which means, that the second term on the r.h.s. of (31) contributes to the vector part of the resulting mass operator (21), while the first term contributes to the scalar part. One should take into account, that in the first term,

$$\Lambda_{\text{scalar}} = \frac{\sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \int_{\tau_{\min}(\mu)}^{\infty} \frac{d\tau \cos(a\sqrt{\tau^2 - 1})}{\sqrt{\tau^2 - 1}} \quad (32)$$

$\tau_{\min}(\mu) = \mu/\sigma r$  for  $\mu > \sigma r$  and 1 otherwise, so that for large  $r, r \gg \frac{\mu}{\sigma}$ , one has the standard  $\mu$ -independent value

$$\Lambda_{\text{scalar}}(r \sim r' > \mu/\sigma) = \frac{\sigma}{\pi^2 r} K_0(a) \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \quad (33)$$

where we have used relations for the McDonald function  $K_0$

$$K_0(a) = \int_0^{\infty} \frac{\cos ax dx}{\sqrt{1 + x^2}}, \quad \int_0^{\infty} da K_0(a) = \frac{\pi}{2}. \quad (34)$$

One can check that at large  $r, r'(r \sim r' > \mu/\sigma)$   $\Lambda_{\text{scalar}} \approx \Lambda_{\text{scalar}}^{(\mu=0)}$  is a smeared  $\delta$ -function,

$$\int \Lambda_{\text{scalar}}^{(\mu=0)}(\mathbf{r}, \mathbf{r}') d^3 \mathbf{r}' = 1. \quad (35)$$

However for large  $\mu, \mu \gg \sqrt{\sigma}$ ,  $\Lambda_{\text{scalar}}$  is different from  $\Lambda_{\text{scalar}}(\mu = 0)$ , and for  $r \sim r' < \mu/\sigma$  one has approximately

$$\Lambda_{\text{scalar}}(\mathbf{r}, \mathbf{r}') = \Lambda_{\text{scalar}}^{(\mu=0)} - f(r, r') \theta(\mu - \sigma r) \quad (36)$$

where

$$f(r, r') = \int_1^{\mu/\sigma r} \frac{d\tau \cos(a\sqrt{\tau^2 - 1})}{\sqrt{\tau^2 - 1}} = \int_1^{\lambda_0} \frac{d\lambda}{\sqrt{1 + \lambda^2}} \cos a\lambda, \quad (37)$$

and  $\lambda_0 = \sqrt{\left(\frac{\mu}{\sigma r}\right)^2 - 1}$ .

Let us now turn to the vector part of interactions,  $\Lambda_{vector}$ ,

$$\begin{aligned} \Lambda_{vector} = & -\frac{2\sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \int_1^{\mu/\sigma r} \frac{\tau d\tau}{\sqrt{\tau^2 - 1}} \cos(a\sqrt{\tau^2 - 1}) \theta(\mu - \sigma r) = \\ & -\hat{1} \frac{2\sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \frac{\sin(a\sqrt{\left(\frac{\mu}{\sigma r}\right)^2 - 1})}{a}. \end{aligned} \quad (38)$$

Returning back to Eq. (21) one can deduce, that

$$M(p_4 = 0, \mathbf{r}, \mathbf{r}') = \sqrt{\pi} T_g J^E(\mathbf{r}, \mathbf{r}') [\Lambda_{scal}(\mathbf{r}, \mathbf{r}') + \gamma_4 \Lambda_{vector}] \equiv M_{scal} + \gamma_4 M_{vect}. \quad (39)$$

Taking into account, that at  $r, r' \gg T_g$  and for the Gaussian  $D^E(x)$  one has from (7)

$$J^E(\mathbf{r}, \mathbf{r}') \cong \frac{(\mathbf{r}\mathbf{r}')}{rr'} 2T_g \sqrt{\pi} D(0) \min(r, r') \quad (40)$$

one has for  $M$  at  $r, r' \gg T_g$  and for  $r \cong r'$

$$M_{scal}(r, r') = \sigma r (\tilde{\delta}^{(3)}(\mathbf{r}, \mathbf{r}') - \xi(\mathbf{r}, \mathbf{r}')) \quad (41)$$

where  $\tilde{\delta}^{(3)}(\mathbf{r}, \mathbf{r}') \equiv \Lambda_{scal}^{(\mu=0)}(\mathbf{r}, \mathbf{r}')$  and

$$\xi(\mathbf{r}, \mathbf{r}') = \delta_\mu = \frac{\sigma}{\pi^2 r} (1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) f(r, r') \theta(\mu - \sigma r) \quad (42)$$

and  $f(r, r')$  is defined in (37).

For  $M_{vect}$  one has, using (38),

$$M_{vect}(\mathbf{r}, \mathbf{r}') = -2\sigma r \varphi_\mu(\mathbf{r}, \mathbf{r}') \theta(\mu - \sigma r) \quad (43)$$

where

$$\varphi_\mu(\mathbf{r}, \mathbf{r}') = \frac{\sigma}{\pi^2 r} \delta(1 - \cos \theta_{\mathbf{r}\mathbf{r}'} ) \frac{\sin(a\sqrt{\left(\frac{\mu}{\sigma r}\right)^2 - 1})}{a}. \quad (44)$$

Note, that one should actually symmetrize all these expressions, e.g.  $\frac{1}{r} \rightarrow \frac{1}{\sqrt{rr'}}$  etc., but we always are in the regime, where  $r \approx r'$ . To estimate the magnitude of nonlocal kernel  $M_{scal}(\mathbf{r}, \mathbf{r}')$  and  $M_{vect}(\mathbf{r}, \mathbf{r}')$  it is convenient to introduce in (18) the local limit of the mass operator, namely

$$\bar{M}_{scal, vect}(r) = \int d^3 \mathbf{r}' M_{scal, vect}(\mathbf{r}, \mathbf{r}'). \quad (45)$$

Exploiting the equalities

$$\int d^3 \mathbf{r}' \tilde{\delta}^{(3)}(\mathbf{r}, \mathbf{r}') = \int d^3 \mathbf{r}' \xi(\mathbf{r}, \mathbf{r}') = \int d^3 \mathbf{r}' \varphi_\mu(\mathbf{r}, \mathbf{r}') = 1 \quad (46)$$

one arrives at the expressions

$$\bar{M}_{scal}(r) = \sigma r \theta(\sigma r - \mu), \quad \bar{M}_{vect} = -2\sigma r \theta(\mu - \sigma r). \quad (47)$$

In the next section we shall discuss approximations made in deriving (41), (43), (47) and physical implications of these results.

## 4 Multiquark states and baryons at nonzero density

Using the local limit for interactions of a light quark with string junction, Eq. (47), one can calculate the change of masses of white states due to density. At this point it is necessary to define exactly the physical meaning of the parameter  $\mu$ , introduced in (10) as the quark chemical potential. That would be true, in the deconfined phase, when quarks can move freely over the volume and occupy the Fermi sphere in momentum space. In the confined phase, however, the term  $\mu$  is meant to be attached to the common string junction of all quark states, constituting the complete set of states (24). This meaning is especially clear from Eq.(21), where  $\mu$  exemplifies the boundary of the occupied states  $\{\varepsilon\}$  in the  $\{n, j, \kappa\}$  notation.

Therefore, assuming that multiquark states located at different string junction points do not overlap, one can associate  $\mu$  for a given quark with the largest value of  $\varepsilon_n$ , occupied by other quarks, belonging to the same string junction point. In what follows we shall show, that in some situations it will be advantageous for two or more nucleons (with different string junctions) to coalesce into a common state with one string junction, building in this way the contracted potential (47).

Note, that for one nucleon (3 quarks) the contraction mechanism does not work and  $\mu$  can be taken at zero value, since no forbidden states exist for each quark of a given color. This, again, is true, provided one-quark states do not overlap at the given density. The situation changes, however, for two nucleons, since e.g. one can add to the first nucleon, i.e. 3 quarks in S-states, 3 quarks in P-states relative to the same string junction point. The latter will move in the potential (47), where  $\mu$  is equal to the S-state Dirac eigenvalue  $\varepsilon_S$ , and the energy  $\varepsilon_P(\mu = \varepsilon_S)$  may become less than  $\varepsilon_S(\mu = 0)$ :

$$\varepsilon_P(\mu = \varepsilon_S) < \varepsilon_S(\mu = 0) \quad (48)$$

The same type of inequality may occur for higher  $L$  states, when more nucleons coalesce to the same Multiquark State (MQS).

This would imply instability of nuclei with respect to transition to MQS matter.

If, however, Eq.(48) does not hold in the limit of zero density, one may have an interesting situation for increasing density, large baryonic chemical potential  $\mu_B$ . Note, that  $\mu_B \neq 3\mu$ , since  $\mu$  is attributed to MQS string junction and does not grow with  $\mu_B$  unless different MQS start to overlap. Hence for  $\mu_B > 0$  (48) should be replaced by

$$\varepsilon_P(\mu = \varepsilon_S) < \varepsilon_S(\mu = 0) + \varepsilon_B(\text{Fermi}). \quad (49)$$

To estimate, let's take  $\varepsilon_S(\mu = 0) \approx 500$  MeV,  $\varepsilon_P(\mu = 500 \text{ MeV}) \approx 630$  MeV (see the Tables 1,2 below), so for  $\varepsilon_B(\text{Fermi}) \approx 130$  MeV the phase transition can occur (which corresponds to densities  $3 \div 4$  times higher than a nuclear one).

Below we perform calculations of  $\varepsilon(\mu)$  for different values of  $\mu$  and  $L = 0, 1$ . We start with the values of  $\varepsilon_{n,L,\kappa}(0)$ , which were computed earlier numerically and analytically using WKB approximation [10] (see Table 1).

In Table 2 we show the change of  $\varepsilon_{nL\kappa}$  as  $\Delta M_S = 3\Delta\varepsilon_n(S)$  for  $S$  and  $\Delta M_P(3q) = 2\Delta\varepsilon_n(S) + \Delta\varepsilon_n(P)$  for the P-wave baryon due to nonzero value of  $\mu$ ,  $\Delta\varepsilon_{nL\kappa}(\mu) = \varepsilon_{nL\kappa}(\mu) - \varepsilon_{nL\kappa}(0)$ , calculated in perturbation theory, considering change of potential in (47) as a



Table 1: Eigenvalues  $\frac{\varepsilon_n}{\sqrt{\sigma}}$  computed numerically, as in (24)

| 1   |               | 0    | 1    | 2    |
|-----|---------------|------|------|------|
| $n$ | sign $\kappa$ |      |      |      |
| 0   | —             | 1.29 | 1.93 | 2.40 |
| 0   | +             | -    | 2.02 | 2.51 |
| 1   | —             | 2.34 | 2.76 | 3.11 |
| 1   | +             | -    | 2.79 | 3.18 |

Table 2: Mass shifts of  $3q$  system due to nonzero  $\mu_q$

| $\mu_q$ (MeV) | $\Delta M_S(3q)$ MeV | $\Delta M_P(3q)$ MeV |
|---------------|----------------------|----------------------|
| 100           | -5                   | -4                   |
| 200           | -70                  | -50                  |
| 300           | -266                 | -200                 |
| 400           | -605                 | -483                 |
| 500           | -1023                | -876                 |

perturbative parameter. One should note at this point, that the vector part of interaction written in (47) for the case of light quark with heavy antiquark, and for the case of 3 quarks interacting with string junction the vector part is transformed as an addition to the color Coulomb potential with coefficient 1/2 (see [11] for explicit derivation of scalar and vector interaction in the nucleon). This prescription was used for calculations in Table 2.


## 5 Discussion of results

Results of the previous section Eqs. (41), (43), (47), can be formulated as follows. The relativistic WKB analysis leads to the  $\mu$ -dependent modification of the confining string, where the piece  $[0, \mu/\sigma]$  of the string near the origin of the string (situated at the heavy quark position in case of heavy-light quark, or at the string junction position in the case of baryons), is dissolved, and the linear confinement starts beyond the critical radius  $r_{cr} = \mu/\sigma$ . Moreover, an attractive vector interaction appears in the same interval with the average magnitude  $\langle \bar{M}_{vect} \rangle \sim \mu$ .

As shown in Table 2 the change in quark energies  $\varepsilon_S(\mu)$ ,  $\varepsilon_P(\mu)$  is large and  $\Delta\varepsilon_P(\mu = 400 \text{ MeV}) = -80 \text{ MeV}$ ,  $\Delta\varepsilon_P(\mu = 500 \text{ MeV}) = -190 \text{ MeV}$ , while  $|\varepsilon_P(\mu = 0) - \varepsilon_S(\mu = 0)| \approx 320 \text{ MeV}$ . Hence it seems to be not advantageous for a quark in neighboring nucleon to add in the P-wave to the  $3q$  system in  $S$  state, with a reduced confinement.

However, the gap is not large. Indeed, the MQS with five quarks in  $S$  state and one quark in  $P$  state will have the mass shift  $\Delta$  only 130 MeV above the NN mass. Note, that phenomenological value of the  $6q$  MQS  $\Delta \cong 210 \text{ MeV}$  is in the same ballpark [14]. Hence

Figure 1: Interactions  $\bar{M}_{scal}(r)$  (solid line) and  $\bar{M}_{vect}(r)$  (dashed line) as functions of distance  $r$ , with  $b = \mu/\sigma$ . Dotted line and dashed line show the qualitative smoothed form of both terms respectively.



for higher density this value of 130 MeV can be compensated by baryon Fermi energy, and the baryon matter can become unstable for transition into MQS matter.

These conclusions should be taken as qualitative. First of all, the WKB method is not a good approximation at small distances, and we have omitted exponentially damped part of  $\psi_n(\mathbf{z})$  in the spectrum, therefore the inner part of the string is to some extent delocalized (see [10] for details) and smoothed, as shown in Fig.1.

Secondly, the interaction between MQS and baryons was not considered above.

Thirdly, we have not taken into account a possible modification and destruction of the vacuum due to the influence of high density quark matter, which might decrease  $\sigma$  or cancel the string completely (as it happens in the thermal phase transition [6]). We have calculated in [5] the critical value of  $\mu_q(T = 0)$ , where deconfinement due to density occurs  $\mu_q^{crit} = \frac{V_1(T^{(0)})}{2} + \left(\frac{9\pi^2}{16n_f}G_2\right)^{1/4} \cong 0.6$  GeV, which corresponds to 7 times normal nuclear density. We are interested here however in the confinement region and small temperatures, with density few times larger than normal.

If however, no density induced vacuum deconstruction takes place, then the resulting physical picture according to Eqs. (47), is the net decreasing of confinement in the inner region of some ensemble of quarks, and appearance of attractive vector potential of the order of  $\mu$  acting on each quark. This may cause creation of partly deconfined bubbles consisting of  $3n$  quarks,  $n = 2, 3, \dots$  in the midst of the nuclear medium, and dynamically is similar to the  $3nq$  bag formation, which was studied before in the framework of the Quark Compound Bag model [14]. Note, however, that bag boundary conditions might be strongly modified as compared to the standard MIT bag model. As it is, preliminary estimates in the previous section demonstrate a possible formation of a new phase of MQS at rather high densities (in heavy ion collisions or in neutron stars) when neutron matter density exceeds 3-4 normal values. A quantitative analysis of this situation needs

a more accurate consideration of the  $3n$  quark system using nonlinear equations for the  $3nq$  Green's function, generalizing Eq. (14).

The formation of these high-density  $3nq$  bubbles may be connected with the explanation so-called cumulative effects in the hadron-nucleus (and nucleus-nucleus) collisions, [15] for an example of this discussion see [16] and refs. therein.

The authors are grateful to N.O.Agasian and A.B.Kaidalov for useful discussions.

This work was supported by the Russian State Corp. "Rosatom", by the President Grant No. NSh-4961.2008.2 for the leading scientific schools, and by the grants RFFI-08-02-00657 and RFFI-09-02-00629. One of the authors (M. A. T.) would like to acknowledge the partial support from the President Grant No. MK-2130.2008.2.

## References

- [1] G. F. Chapline, M. H. Johnson, E. Teller, M. S. Weiss, Phys. Rev. **D8** (1973) 4302; T. D. Lee, G. C. Wick, Phys. Rev. **D9** (1974) 2291; T. D. Lee, Rev. Mod. Phys. **47** (1975) 267.
- [2] M. Baldo and C. Maieron, J. Phys **G34** (2007) R243 [arXiv:nucl-th/0703004].
- [3] Yu. A. Simonov, JETP Lett. **54** (1991) 249; **55** (1992) 605; Phys. At. Nucl. **58** (1995) 309.
- [4] M. D'Elia, A. Di Giacomo, E. Meggiolaro, Phys. Lett. **B408**, (1997) 315; Phys. Rev **D67** (2003) 114504.; A. Di Giacomo, E. Meggiolaro, H. Panagopoulos, Nucl. Phys. **B483** (1997) 371.
- [5] Yu.A.Simonov, M.A.Trusov, Phys. Lett. D **650** (2007) 36; JETP Lett. **85** (2007) 730.
- [6] A.V.Nefediev, Yu.A.Simonov, M.A.Trusov, Int. J. Mod. Phys. E **18** (2009) 1.
- [7] Yu.A.Simonov, Phys. Atom. Nucl. **609** (1997) 2069.
- [8] P. J. Ellis, J. I. Kapusta and H. B. Tang, Phys. Lett. **B443** (1998) 63; N. O. Agasian, Phys. Lett. **B519** (2001) 71; JETP Lett. **74** (2001) 353; N. O. Agasian and S. M. Fedorov, JETP Lett. **78** (2003) 607; N. O. Agasian, JETP Lett. **57** (1993) 208.
- [9] S. V. Ivanov, G. P. Korchemsky, Phys. Lett. **B154** (1985) 197; S. V. Ivanov, G. P. Korchemsky, A. V. Radyushkin, Sov.Y. Nucl. Phys. **44** (1986) 145; V. I. Shevchenko and Yu. A. Simonov, Phys. Lett. **B437** (1998) 146.
- [10] Yu. A. Simonov, Phys. At. Nucl. **60** (1997) 2069; arXiv:hep-ph/9704301; Yu. A. Simonov, Phys. Rev. **D65** (2002) 094018; arXiv:hep-ph/0201170.
- [11] V. S. Popov, D. V. Voskresensky, V. L. Eletsky, V. D. Mur, JhETF **76** (1979) 431; V. D. Mur, V. S. Popov, Yad.Fiz. **28** (1978) 837; V. D. Mur, V. S. Popov, D. V. Voskresensky, Pis'ma JhETF **28** (1978) 140.

- [12] V.D.Mur, V.S.Popov, Yu.A.Simonov and V.P.Yurov, JETP (1994) 1; Yu.A.Simonov, J.A.Tjon, Phys. Rev. **62** (2000) 014501.
- [13] Yu.A.Simonov, Phys. Atom. Nucl. **66** (2003) 338; arXiv: hep-ph/0205334;
- [14] Yu. A. Simonov, Nucl. Phys. **A416** (1984) 109; **A463** (1987) 231;  
Yu. S. Kalashnikova, I. M. Narodetskii, Int. J. Mod. Phys. **A4** (1989) 335.
- [15] Yu.D.Bayukov et al., Izvestia Akad. Nauk SSSR, Fizika, **30** (1966) 521; Yad. Fiz **18** (1973) 1246; A.M.Baldin, Short Communications in Physics, Lebedev Inst., 1 (1971) 35.
- [16] A. V. Efremov, A. B. Kaidalov, et al. Yad Fiz. **47** (1988) 1364.